

QUANTUM CARPETS MADE SIMPLE¹

I. Marzoli, F. Saif, I. Bialynicki-Birula*, O. M. Friesch, A. E. Kaplan[†],
W. P. Schleich²

Abteilung für Quantenphysik, Universität Ulm, D-89069 Ulm, Germany

**Center for Theoretical Physics, Lotników 46, 02-668 Warsaw, Poland*

[†]Electr. & Comp. Eng. Dept., The Johns Hopkins University, MD-21210, USA

We show that the concept of degeneracy is the key idea for understanding the quantum carpet woven by a particle in the box.

1 Introduction

Interesting structures [1, 2] emerge in the space-time representation of the probability distribution for a particle in the box, as shown in Fig. 1. Three explanations of these quantum carpets offer themselves: Interference terms in the Wigner function [3, 4], degeneracy of intermode traces [5, 6] and cancelation between appropriate terms of the energy representation [2, 7, 8]. All of these explanations are rather involved. We therefore in the present paper develop a simple argument for this surprising phenomenon.


We identify three properties of the particle in the box as the thread of the quantum carpets:

- (i) The probability density involves the product of two standing waves creating contributions with the sum and the difference of the wave numbers.
- (ii) The quadratic dispersion relation connecting the energy and the momentum gives rise to a multi-degeneracy.
- (iii) The appropriate initial conditions enhance this degeneracy.

The paper is organized as follows. In Sect. 2 we briefly review the important formulas of the problem of the particle in the box. We then in Sect. 3 give a heuristic argument for the quantum structures. In the Appendix we derive a summation formula which allows us in Sect. 4 to cast the probability density into a form which brings out most clearly the canals and ridges of the quantum carpets. We conclude by summarizing the main results in Sect. 5.

¹Special Issue on Quantum Optics and Quantum Information

²E-mail address: schleich@physik.uni-ulm.de



Quantum carpet woven by a particle moving with a non-relativistic velocity in a one-dimensional box. The carpet arises from the space-time representation of the probability density. As initial condition of the Schrödinger equation with the non-relativistic Hamiltonian $H_{nr} = p^2/(2M)$ we have used a Gaussian wave packet of width $\Delta x = L/20$ and average wave number $\bar{k} = 10\pi/L$ located at $\bar{x} = L/2$.

2 The particle in the box: Fundamentals

In the present section we summarize the essential ingredients of the problem of the particle in the box. In particular, we focus on the energy representation of the wave function.

The probability amplitude $\psi(x, t)$ to find the particle of mass M at time t at the position x in the box of length L reads

$$\psi(x, t) = \sum_{m=1}^{\infty} \psi_m u_m(x) \exp\left(-\frac{i}{\hbar} E_m t\right). \quad (1)$$

Here the quantities

$$\psi_m \equiv \int_0^L dx \varphi(x) u_m(x) \quad (2)$$

are the expansion coefficients of the initial wave packet $\psi(x, t = 0) \equiv \varphi(x)$ into the energy wave functions [9]

$$u_m(x) \equiv \sqrt{\frac{1}{2L}} \frac{1}{i} (e^{ik_m x} - e^{-ik_m x}) \quad (3)$$

with wave numbers

$$k_m \equiv mk_1 \equiv m \frac{\pi}{L} \quad (4)$$

and eigen energies

$$E_m \equiv \frac{(\hbar k_m)^2}{2M} = m^2 E_1 = m^2 \hbar \omega_1 \equiv m^2 \hbar \frac{2\pi}{T}. \quad (5)$$

In the last step we have introduced [10, 11] the revival time

$$T \equiv \frac{4ML^2}{\hbar\pi} \quad (6)$$

where the wave function is identical to its initial form at $t = 0$, that is $\psi(x, t = T) = \psi(x, t = 0)$.

We conclude this summary of the important equations by deriving a representation of the probability amplitude ψ that is slightly different from Eq. (1). In Sect. 4 we will use this expression to bring out the location and shape of the structures in the quantum carpet.

We substitute the energy wave function u_m into Eq. (1) and use the expressions Eqs. (4) and (5) for the wave numbers k_m and energies E_m . We then arrive at

$$\psi(x, t) = \frac{1}{i\sqrt{2L}} \sum_{m=1}^{\infty} \left\{ \psi_m \exp \left[i\pi m \left(\frac{x}{L} - m \frac{2t}{T} \right) \right] - \psi_m \exp \left[-i\pi m \left(\frac{x}{L} + m \frac{2t}{T} \right) \right] \right\}. \quad (7)$$

When we define the expansion coefficients ψ_m for negative values of m by

$$\psi_{-|m|} \equiv -\psi_{|m|} \quad (8)$$

we find the compact representation

$$\psi(x, t) = \frac{1}{i\sqrt{2L}} \sum_{m=-\infty}^{\infty} \psi_m \exp \left[i\pi m \left(\frac{x}{L} - m \frac{2t}{T} \right) \right] \quad (9)$$

of the wave function.

3 Quantum carpets: A heuristic argument

In the present section we use the expression Eq. (1) to show that the probability density consists of four terms: Two terms correspond to the classical trajectories, whereas the other two represent the striking canals and ridges of Fig. 1.

Since the structures appear in the probability density

$$W(x, t) \equiv \psi^*(x, t) \psi(x, t) \quad (10)$$

we now use the energy representation Eq. (1) of ψ to find W and arrive at

$$W(x, t) = -\frac{1}{2L} \sum_{m,n=1}^{\infty} \psi_m^* \psi_n (e^{ik_m x} - e^{-ik_m x})(e^{ik_n x} - e^{-ik_n x}) \exp \left[i(k_m^2 - k_n^2) \frac{\hbar t}{2M} \right]. \quad (11)$$

Here we have used the expressions, Eqs. (3) and (5), for the energy wave functions and the energies.

When we multiply out the individual waves in the product of the two energy wave functions we recognize that the probability

$$W = I_{qc}^{(+)} + I_{qc}^{(-)} + I_{cl}^{(+)} + I_{cl}^{(-)} \quad (12)$$

consists of four contributions. The terms

$$I_{qc}^{(\pm)}(x, t) \equiv -\frac{1}{2L} \sum_{m,n=1}^{\infty} \psi_m^* \psi_n \exp \left\{ \pm i(k_m + k_n) \left[x \pm (k_m - k_n) \frac{\hbar t}{2M} \right] \right\} \quad (13)$$

arise from the multiplication of the two co-propagating waves $\exp(\pm ik_m x)$ and $\exp(\pm ik_n x)$ in the two energy wave functions. Note that the relation

$$k_m^2 - k_n^2 = (k_m + k_n)(k_m - k_n) \quad (14)$$

has allowed us to factor out the sum $k_m + k_n$ of the wave numbers in the expression Eq. (13). This creates the difference $k_m - k_n$ of the wave numbers in the expression in the square brackets.

In contrast the terms

$$I_{cl}^{(\pm)}(x, t) \equiv \frac{1}{2L} \sum_{m,n=1}^{\infty} \psi_m^* \psi_n \exp \left\{ \pm i(k_m - k_n) \left[x \pm (k_m + k_n) \frac{\hbar t}{2M} \right] \right\} \quad (15)$$

are a consequence of the multiplication of the two counter-propagating waves $\exp(\pm ik_m x)$ and $\exp(\mp ik_n x)$ in the two energy wave functions. Here the factorization property Eq. (14) has led to the difference $k_m - k_n$ of the wave numbers outside the square brackets and the sum $k_m + k_n$ inside.

The phases

$$\phi_{m,n}^{(\pm)}(x, t) \equiv \frac{x}{L} \pm (k_m - k_n) \frac{\hbar t}{2ML} = \frac{x}{L} \pm (m - n) \frac{t}{T/2} \quad (16)$$

and

$$\Phi_{m,n}^{(\pm)}(x, t) \equiv \frac{x}{L} \pm (k_m + k_n) \frac{\hbar t}{2ML} = \frac{x}{L} \pm (m + n) \frac{t}{T/2} \quad (17)$$

in the square brackets of Eqs. (13) and (15) correspond to straight lines in space-time. The steepness of these world lines is determined by the difference and the sum of the quantum numbers m and n . This opens the possibility for a multi-degeneracy: Different pairs of quantum numbers m and n can give rise to the same difference $m - n$ or sum

$m + n$. Therefore many world lines can lie on top of each other enhancing in this way the contrast of the structures.

It is the expansion coefficients ψ_m that decide the question of enhancement or suppression of these world lines. To understand this we consider a distribution ψ_m of energy excitations which satisfies the factorization property

$$\psi_m^* \psi_n = \psi_{m+n}^{(+)} \psi_{m-n}^{(-)}. \quad (18)$$

Hence we can replace the product $\psi_m^* \psi_n$ of the initial expansion coefficients by another product of new expansions coefficients $\psi_s^{(+)}$ and $\psi_r^{(-)}$ which now only depend on the sum s and the difference r of the quantum numbers. Any Gaussian wave packet satisfies this condition.

We conclude this section by considering a Gaussian wave packet centered at quantum number \bar{m} and width Δm such that $1 \ll \Delta m \ll \bar{m}$. In this case we find a clear separation of classical and quantum trajectories contributing to the probability density: The terms $I_{cl}^{(\pm)}$ contain the classical trajectories whereas the terms $I_{qc}^{(\pm)}$ are the origin of the carpet. In order to bring this out we recall that the terms $I_{cl}^{(\pm)}$ contain the phase $\Phi_{m,n}^{(\pm)}$ with the sum of the quantum numbers. Due to the Gaussian weight factor with its maximum at $\bar{m} \gg 1$ this gives rise to large quantum numbers of the order of $2\bar{m}$. The steepness of the corresponding world lines is therefore proportional to $\bar{m}^{-1} \ll 1$. Consequently the world lines are rather flat and correspond to the classical trajectories. In contrast the terms $I_{qc}^{(\pm)}$ contain the phase $\phi_{m,n}^{(\pm)}$ with the difference of the quantum numbers. This corresponds to steep world lines—the striking canals and ridges.

4 A new representation of the probability density

In the preceding section we have shown that for the evaluation of $I_{qc}^{(\pm)}$ and $I_{cl}^{(\pm)}$ it is natural to introduce the new summation indices $m \pm n$. In the present section we pursue this idea. However, we do not start from Eqs. (13) and (15) but from Eq. (9).

The probability W then reads

$$W(x, t) = \frac{1}{2L} \sum_{m,n=-\infty}^{\infty} \psi_m^* \psi_n \exp \left\{ -i\pi(m-n) \left[\frac{x}{L} - (m+n) \frac{2t}{T} \right] \right\}. \quad (19)$$

With the help of the summation formula

$$\sum_{m,n=-\infty}^{\infty} f_{m,n} = \frac{1}{2} \sum_{l=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} (-1)^{jl} \int_{-\infty}^{\infty} d\rho f \left[\frac{1}{2}(j+\rho), \frac{1}{2}(j-\rho) \right] \exp(i\pi l\rho) \quad (20)$$

derived in the Appendix the probability density takes the form

$$W(x, t) = \frac{1}{4L} \sum_{l,j=-\infty}^{\infty} (-1)^{jl} \Psi^{(W)} \left[\frac{j}{2}, \chi_{j,l}(x, t) \right]. \quad (21)$$

Here we have introduced the Wigner function

$$\Psi^{(W)}(\mu, \xi) \equiv \int_{-\infty}^{\infty} d\rho \psi^* \left[\mu + \frac{\rho}{2} \right] \psi \left[\mu - \frac{\rho}{2} \right] \exp(-i\pi\rho\xi), \quad (22)$$

of the expansion coefficients ψ_m . Note that $\psi[\mu]$ is a continuous extension of ψ_m such that $\psi[\mu] \equiv \psi_m$ for $\mu = m$.

Moreover, we have defined

$$\chi_{j,l}(x, t) \equiv \frac{x}{L} - j \frac{t}{T/2} - l. \quad (23)$$

This expression brings out most clearly that the probability density consists of a superposition of structures $\Psi^{(W)}$ aligned along straight lines defined by $\chi_{j,l}(x, t)$. The Wigner function $\Psi^{(W)}$ of the expansion coefficients determines the shape of the structures.

We conclude this section by making contact with the expression,

$$W(x, t) = \frac{\pi\hbar}{2L} \sum_{l,j=-\infty}^{\infty} (-1)^{jl} \Psi_{\phi}^{(W)}[\chi_{j,l}(x, t), p_j], \quad (24)$$

derived in Ref. [3]. Here $p_j \equiv j\pi\hbar/(2L)$ and

$$\Psi_{\phi}^{(W)}(x, p) = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dy \phi \left(x - \frac{y}{2} \right) \phi^* \left(x + \frac{y}{2} \right) \exp(-ipy/\hbar) \quad (25)$$

is the Wigner function [12] of the superposition wave function

$$\phi(x) \equiv \varphi(x) - \varphi(-x) \quad (26)$$

built out of the original wave packet and its mirror image.

For this purpose we substitute the representation of the expansion coefficients ψ_m in terms of the original wave packet $\varphi(x)$ in position space Eq. (2) and the energy wave functions u_m Eq. (3) into the definition Eq. (22) of the Wigner function $\Psi^{(W)}$. The integration over ρ leads to a Dirac delta function which allows us to perform one of the integrations over position. The remaining expression is then indeed the Wigner function $\Psi_{\phi}^{(W)}$ of the superposition wave function ϕ .

5 Conclusions

In the energy representation the probability of finding the particle at time t at position x is a double sum over all energy quantum numbers. The quadratic dispersion relation of the free particle allows us to express this double sum by another double sum extending over the sums and differences of the quantum numbers. In this way we represent the probability distribution in space-time as a superposition of structures along straight world lines. Their steepness and their starting point are determined by integers. The

shape of the initial wave packet, that is the form ψ_m of the initial excitation of the energy eigenstates governs the shape of these structures.

We emphasize that the expression Eq. (21) for the probability density derived in the present paper is very different from the one which follows from the representation [2, 10, 11]

$$\psi\left(x, t = \frac{q}{r}T + \Delta t\right) = \sum_{l=-\infty}^{\infty} \mathcal{W}_l^{(r)} \varphi\left(x - \frac{l}{r}2L, \Delta t\right) - \sum_{l=-\infty}^{\infty} \mathcal{W}_l^{(r)} \varphi\left(-x + \frac{l}{r}2L, \Delta t\right), \quad (27)$$

of the wave function ψ in the neighborhood of a fraction q/r of the revival time T . Here $\mathcal{W}_l^{(r)}$ denotes the Gauss sums [13, 14] and

$$\varphi(x, t) \equiv \int_0^L dx' \varphi(x') G_{free}(x, t|x', 0) \quad (28)$$

is the initial wave function $\varphi(x) \equiv \psi(x, t = 0)$ propagated freely according to the Green's function G_{free} of the free particle. Indeed the above formula Eq. (27) is a more local representation in space-time whereas Eq. (21) is a more global one: It depicts the probability density W as a superposition of structures along *vertical* and *tilted* world lines, whereas the revival representation Eq. (27) uses a superposition of structures along lines of constant time, that is along horizontal world lines.

We have obtained these results using the summation formula derived in the Appendix. This formula has a much wider range of application. For example, it immediately provides an expression for the slightly relativistic particle. In this case the structures are not along straight, but curved lines as shown in Fig. 2. Moreover, it provides insight into the development of fractal canals discovered in Ref. [2], when the initial wave packet is constant. However, space does not allow us to go deeper into these topics of future publications.

Appendix: A useful summation formula

In this appendix we derive two different but equivalent representations of the double sum

$$I \equiv \sum_{m,n=-\infty}^{\infty} f_{m,n} \quad (29)$$

with coefficients $f_{m,n}$.

Our derivation relies on the introduction of the new summation indices $m + n \equiv s$ and $m - n \equiv r$. Note however, that this definition puts certain restrictions on s and r . Indeed, we have to distinguish two cases: (i) when m and n are both even or odd, and (ii) when one of them is odd and the other is even. In the case (i) we find that $m - n$ and $m + n$ are both even. Hence we have the substitutions

$$m - n \equiv 2r \quad \text{and} \quad m + n \equiv 2s. \quad (30)$$

Quantum carpet woven by a particle moving with a slightly relativistic velocity in a one-dimensional box. We use the Schrödinger equation to propagate a Gaussian wave packet with the Hamiltonian $H_r \equiv H_{nr}[1 - H_{nr}/(2Mc^2)]$. The straight canals and ridges of the non-relativistic box problem of Fig. 1 are now curved. Here we have chosen the initial conditions

In the case (ii) we find that $m - n$ and $m + n$ are both odd, which leads to the definition

$$m - n \equiv 2r + 1 \quad \text{and} \quad m + n \equiv 2s + 1. \quad (31)$$

We therefore find the rule

$$\sum_{m,n=-\infty}^{\infty} f_{m,n} = \sum_{r,s=-\infty}^{\infty} f_{s+r,s-r} + \sum_{r,s=-\infty}^{\infty} f_{s+r+1,s-r} \quad (32)$$

for replacing the original sums extending over m and n by new sums extending over r and s .

We can combine the two terms in Eq. (32) into one, when we replace either the summation over r , or the one over s , by an integration. The Poisson summation formula [15]

$$\sum_{m=-\infty}^{\infty} g_m = \sum_{l=-\infty}^{\infty} \int_{-\infty}^{\infty} d\mu g[\mu] \exp(2\pi il\mu) \quad (33)$$

allows us to do this in an exact way. Here $g[\mu]$ is an extension of the function g_m to the whole real axis such that $g[\mu]$ takes on the values g_m at integer values $\mu = m$.

When we apply the Poisson formula to the summation over r we arrive at

$$I = \sum_{l=-\infty}^{\infty} \left\{ \sum_{s=-\infty}^{\infty} \int_{-\infty}^{\infty} d\rho f[s + \rho, s - \rho] \exp(2\pi i l \rho) + \sum_{s=-\infty}^{\infty} \int_{-\infty}^{\infty} d\rho f[s + \rho + 1, s - \rho] \exp(2\pi i l \rho) \right\}, \quad (34)$$

which after the substitutions $\tilde{\rho} \equiv 2\rho$ and $\bar{\rho} \equiv 2\rho + 1$ in the two integrals takes the form

$$I = \frac{1}{2} \sum_{l=-\infty}^{\infty} \left\{ \sum_{s=-\infty}^{\infty} \int_{-\infty}^{\infty} d\tilde{\rho} f \left[\frac{1}{2}(2s + \tilde{\rho}), \frac{1}{2}(2s - \tilde{\rho}) \right] \exp(i\pi l \tilde{\rho}) + \sum_{s=-\infty}^{\infty} (-1)^l \int_{-\infty}^{\infty} d\bar{\rho} f \left[\frac{1}{2}(2s + 1 + \bar{\rho}), \frac{1}{2}(2s + 1 - \bar{\rho}) \right] \exp(i\pi l \bar{\rho}) \right\}. \quad (35)$$

Here we have used for the last integral the relation

$$\exp(i\pi l) = (-1)^l, \quad (36)$$

and have written the arguments of the function f in a way that brings out most clearly that the two integrals are the even and odd terms of a single sum. The only obstacle left before we can combine these two terms is the term $(-1)^l$. When we recall that

$$(-1)^{jl} = \begin{cases} (-1)^{2sl} = 1 & \text{for } j = 2s \\ (-1)^{(2s+1)l} = (-1)^l & \text{for } j = 2s + 1 \end{cases} \quad (37)$$

we find indeed

$$\sum_{m,n=-\infty}^{\infty} f_{m,n} = \frac{1}{2} \sum_{l=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} (-1)^{jl} \int_{-\infty}^{\infty} d\rho f \left[\frac{1}{2}(j + \rho), \frac{1}{2}(j - \rho) \right] \exp(i\pi l \rho). \quad (38)$$

We conclude this appendix by presenting a different expression for the double sum I which follows from Eq. (32) when we replace the summation over s by an integration using the Poisson summation formula, Eq. (33). In this case we find following the same train of thought

$$\sum_{m,n=-\infty}^{\infty} f_{m,n} = \frac{1}{2} \sum_{l=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} (-1)^{jl} \int_{-\infty}^{\infty} d\sigma f \left[\frac{1}{2}(\sigma + j), \frac{1}{2}(\sigma - j) \right] \exp(i\pi l \sigma). \quad (39)$$

We recognize that the two representations are different: In the one of Eq. (38) the integration variable ρ enters in an asymmetric way whereas in the one of Eq. (39) the integration variable σ appears in a symmetric way. Nevertheless, both representations are completely equivalent.

Acknowledgement

We express our gratitude to P. J. Bardroff, M. V. Berry, J. H. Eberly, M. Fontenelle, F. Großmann, M. Hall, H. J. Kimble, T. Kiss, W. E. Lamb, Jr., K. A. H. van Leeuwen, C. Leichtle, J. Marklof, M. M. Nieto, J. M. Rost, E. C. G. Sudarshan and P. Stifter for many fruitful discussions on this topic. Two of us (I. B.-B.) and (A. E. K.) thank the Humboldt Stiftung for their support.

References

- [1] W. Kinzel: *Phys. Bl.* **51** (1995) 1190;
- [2] M. V. Berry: *J. Phys. A* **29** (1996) 6617; M. V. Berry, S. Klein: *J. Mod. Optics* **43** (1996) 2139;
- [3] P. Stifter, C. Leichtle, W. P. Schleich, J. Marklof: *Z. Naturf.* **52 a** (1997) 377;
- [4] I. Marzoli, O. M. Friesch, W. P. Schleich: *Proceedings of the Fifth Wigner Symposium*, ed. P. Kasperkovitz, World Scientific, in press;
- [5] A. E. Kaplan, P. Stifter, K. A. H. van Leeuwen, W. E. Lamb, Jr., W. P. Schleich: *Physica Scripta* (in press) ;
- [6] I. Marzoli, I. Bialynicki-Birula, O. M. Friesch, A. E. Kaplan, W. P. Schleich: *Proceedings of the Physical Research Laboratory Jubilee Conference on Nonlinear Dynamics and Computational Physics*, ed. V. B. Sheorey, Narosa Publishing House, in press [see also *Los-Alamos e-print archive* quant-ph/9804015 (1998)];
- [7] F. Großmann, J.-M. Rost, W. P. Schleich: *J. Phys. A* **30** (1997) L277;
- [8] J. Marklof: *Limit theorems for theta sums with applications in quantum mechanics* Shaker Verlag, 1997;
- [9] See, e.g. D. Bohm: *Quantum theory*, Prentice-Hall, 1951;
- [10] P. Stifter, W. E. Lamb, Jr., W. P. Schleich: *Proceedings of the Conference on Quantum Optics and Laser Physics*, ed. L. Jin and Y. S. Zhu, World Scientific, 1997; I. Marzoli, O. M. Friesch, P. Stifter, W. P. Schleich: to be published;
- [11] D. L. Aronstein, C. R. Stroud, Jr.: *Phys. Rev. A* **55** (1997) 4526;
- [12] M. Hillery, R. F. O'Connell, M. O. Scully, E. P. Wigner: *Phys. Rep.* **106** (1984) 121;
- [13] S. Lang: *Algebraic Number Theory* Addison-Wesley, 1970;
- [14] J. H. Hannay, M. V. Berry: *Physica D* **1** (1980) 267;
- [15] M. J. Lighthill: *Introduction to Fourier analysis and generalized functions*, Cambridge University Press, 1960.